

THE CLUSTER $L_X - \sigma$ RELATION HAS IMPLICATIONS FOR SCALE-FREE COSMOLOGIES

ANDISHEH MAHDAVI¹

Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138

Submitted March 24, 1999; Accepted May 20, 1999 for publication in The Astrophysical Journal Letters

ABSTRACT

I show that the cluster $L_X - \sigma$ relation should be sensitive to cosmologies with a scale-free power spectrum of initial density fluctuations, $P(k) \propto k^n$. I derive the dependence, and argue that a conservative interpretation of current observations implies $n < -2.0$ and $n < -1.1$ at the one-sided 90% and 99% confidence levels, respectively. This result, which agrees with constraints on n from the x-ray cluster temperature function, should be roughly independent of the value of Ω or Λ .

Subject headings: Galaxies: clusters: general — cosmology: theory — X-rays: galaxies

1. INTRODUCTION

The dynamical state of a system of galaxies may be characterized by three intimately related physical quantities: the bolometric x-ray luminosity L_X , the average emission-weighted plasma temperature $\langle T \rangle$, and the average dark matter velocity dispersion, σ_{dm} (as traced by the projected velocity dispersion of the galaxies, σ_p). There is an observed correlation among these parameters, crudely given by $\langle T \rangle \propto \sigma_p^{\alpha_1}$, $L_X \propto \langle T \rangle^{\alpha_2}$, and $L_X \propto \sigma_p^{\alpha_3}$, and roughly in agreement with the predictions of physical models. My goal in this letter is to show that the value of the slope α_3 at $z = 0$ can constrain the spectrum of the primordial density fluctuations, $P(k) \propto k^n$.

One powerful tool for linking the x-ray properties of galaxy systems with the cosmological parameters already exists. The cluster temperature function, which is directly related to the cluster mass distribution, can be used at zero redshift to measure n . Henry & Arnaud (1991) analyze data from the Einstein satellite to obtain $n \approx -1.7 \pm 0.55$; data from the ROSAT and ASCA missions (Markevitch 1998), corrected for the presence of cooling flows, implies a somewhat steeper $n \approx -2 \pm 0.3$. The redshift evolution of the cluster temperature function can break the degeneracy between the density parameter, Ω , and the normalization of the primordial spectrum (Henry 1997).

Here I introduce a new zero-redshift method. I suggest that α_3 , the slope of the $L_X - \sigma_p$ relation, is related to n in a way that is complementary to the dependence of the temperature function on n . While it is important to have a flux-limited sample for the temperature function method, using α_3 to compute n requires only a large number of pointed observations of systems of galaxies. In §2 I derive the relationship between α_3 and n . In §3 I compare the results with available observations, and discuss possible systematic biases. In §4 I summarize.

2. DERIVATION

Here I use some of the definitions in the paper by Navarro, Frenk, & White (1997; NFW). My results, however, are largely independent of their N-body simulations of dark matter halos, and apply to density profiles different from the one they introduce. Eke, Navarro, & Frenk

(1998) provide a derivation of the $L_X - \langle T \rangle$ relation which is analogous to part of what follows, but does not focus on any cosmological use of α_2 .

2.1. Halo Parameters and Definitions

A spherically symmetric dark matter halo may be characterized by r_{200} , the radius which encloses 200 times the critical density of the universe, ρ_{crit} . The characteristic mass, M_{200} , is then

$$M_{200} = \frac{800\pi}{3} r_{200}^3 \rho_{crit} \quad (1)$$

$$\rho_{crit} = \frac{3H_0^2}{8\pi G} Z(z), \quad (2)$$

$$Z(z, \Omega) = (1+z)^3 \frac{\Omega_0}{\Omega(z)}. \quad (3)$$

Here H_0 is the Hubble constant, Ω is the density parameter, and z is the redshift. The halo's characteristic circular velocity is $V_{200} = \sqrt{GM_{200}/r_{200}}$, or, with the substitution of equation (1),

$$V_{200} = 10H_0 r_{200} Z^{1/2}. \quad (4)$$

Now consider a halo density profile of the form,

$$\rho_{dm}(r) = \rho_{crit} \delta_c \tilde{\rho}\left(\frac{cr}{r_{200}}\right), \quad (5)$$

where c is the halo concentration, δ_c is the characteristic density, and $\tilde{\rho}(y)$ is a nonnegative, declining function of y . Then equation (1) requires the following relation between δ_c and c ,

$$\delta_c = \frac{200c^3}{3C(c)}, \quad (6)$$

$$C(y) = \int_0^y t^2 \tilde{\rho}(t) dt. \quad (7)$$

In scale-free cosmologies, the characteristic density also depends on n , the slope of the initial fluctuation spectrum, because c should trace the mean density of the universe at the time of the halo's formation. NFW use Press-Schechter

¹amahdavi@cfa.harvard.edu

theory to examine the exact relationship among δ_c , M_{200} , and n . They find that the characteristic density should be proportional to the mean density of the universe at the epoch when the nonlinear mass M_* is (1) a fixed and (2) a very small fraction of the current halo mass. This implies that δ_c should scale with M and n the same way as the average background density scales with M_* and n :

$$\delta_c \propto M_{200}^{-(n+3)/2}. \quad (8)$$

The slope of this scaling relation is better than 5% accurate for all $\Omega \leq 1$ cosmologies with either $\Lambda = 0$ or $\Omega + \Lambda = 1$, where Λ is the cosmological constant in units of $3H^2$. With the substitution $\gamma = -(n+3)/2$ and use of equations (1), (4), and (6),

$$\frac{c^3}{C(c)} \propto V_{200}^{3\gamma} Z^{-\gamma/2}. \quad (9)$$

Next, I compute the circular velocity profile of the halo, $V_{circ} = \sqrt{GM(r)/r}$. Under the transformation $r \rightarrow yr_{200}/c$,

$$V_{circ}(y)^2 = V_{200}^2 \frac{cC(y)}{yC(c)} \quad (10)$$

Note that the dark matter velocity dispersion profile, $\sigma_{dm}(y)$, will not in general have the same shape as $V_{circ}(y)$, contrary to the assumption in similar derivations (e.g. Eke et al. 1998). Rather, the velocity dispersion of the halo is given by the Jeans equation for a spherical, nonrotating system of collisionless particles (Binney & Tremaine 1987):

$$\frac{d(\sigma_r^2 \rho)}{dr} + 2\beta_J \rho \sigma_r^2 = -\frac{GM\rho}{r^2}. \quad (11)$$

Here σ_r is the radial velocity dispersion, β_J is the velocity anisotropy parameter, and M is the mass inside the radius r . For systems with isotropic velocity dispersion tensors, $\beta_J = 0$, and locally $\sigma_{dm}^2 = 3\sigma_r^2$. In terms of the previously defined quantities, equation (11) has the solution

$$\sigma_r^2(y) = V_{200}^2 \frac{c}{C(c)} \tilde{\sigma}^2(y), \quad (12)$$

$$\tilde{\sigma}^2(y) = \frac{1}{\tilde{\rho}(y)} \int_y^\infty \frac{\tilde{\rho}(t)C(t)}{t^2} dt, \quad (13)$$

Finally, consider the average dark matter velocity dispersion, $\langle \sigma_{dm}^2 \rangle$, within the virial radius r_{200} :

$$\langle \sigma_{dm}^2 \rangle = \frac{\int_0^c 3\sigma_r^2(y) \tilde{\rho}(y) y^2 dy}{\int_0^c \tilde{\rho}(y) y^2 dy} \quad (14)$$

$$= V_{200}^2 \frac{cD(c)}{C(c)^2}, \quad (15)$$

$$D(y) = \int_0^y 3\tilde{\sigma}^2(t) \tilde{\rho}(t) t^2 dt. \quad (16)$$

2.2. X-Ray Luminosity

The x-ray luminosity of a ball of plasma is

$$L_X = \int \lambda(T) n_e n_i dV. \quad (17)$$

Here $\lambda(T)$ is the bolometric emissivity as a function of the local temperature T , n_e is the electron number density, n_i is the ion number density, and the integral is over the entire volume of the system. Now if (1) $n_i = n_e$, (2) the system is spherically symmetric, and (3) the gas density is related to the dark matter distribution by $\rho_{gas}(r) = f\rho_{dm}(r)$, where f is the gas mass fraction, then

$$L_X = 4\pi \int_0^\infty \lambda(T) \left(\frac{f\rho_{dm}}{\mu m_p} \right)^2 r^2 dr, \quad (18)$$

where $\mu \equiv \rho_{gas}/(n_i m_p)$ is the mean molecular weight. Substitution of equation (5) yields,

$$L_X = 4\pi \left(\frac{f\rho_{crit}}{\mu m_p} \right)^2 \frac{\delta_c^2}{c^3} r_{200}^3 \int_0^\infty \lambda(T) \tilde{\rho}(y)^2 y^2 dy. \quad (19)$$

Now eliminate r_{200} , δ_c , and ρ_{crit} using equations (2), (4) and (6):

$$L_X = \frac{5H_0 Z^{1/2}}{2\pi G^2} \left(\frac{f}{\mu m_p} \right)^2 \frac{c^3}{C(c)^2} V_{200}^3 \times \int_0^\infty \lambda(T) \tilde{\rho}(y)^2 y^2 dy. \quad (20)$$

Next, note that thermal bremsstrahlung is the dominant cooling process for rich clusters of galaxies; hence $\lambda(T) \propto T^{1/2}$. Then, if the gas is in local hydrostatic equilibrium with the dark matter, $T(r) \propto \sigma_{dm}(r)^2$, and hence $\lambda(T, r) \propto \sigma_{dm}(r)$. Leaving out the constants and substituting equation (12) yields,

$$L_X \propto f^2 Z^{1/2} \frac{c^{7/2}}{C(c)^{5/2}} V_{200}^4 \int_0^\infty 3\tilde{\sigma}(y) \tilde{\rho}(y)^2 y^2 dy \quad (21)$$

The integral on the right-hand side is just a number, and may be dropped.

Now one must replace V_{200} , which is not observable, with the projected galaxy velocity dispersion σ_p , which can be determined from optical surveys. In the cores of relaxed clusters σ_p^2 is proportional to $\langle \sigma_{dm}^2 \rangle$, the average dark matter velocity dispersion (equation 14). Then

$$L_X \propto f^2 Z^{1/2} \frac{c^{3/2} C(c)^{3/2}}{D(c)^2} \sigma_p^4. \quad (22)$$

If f and c remain constant, the above expression reduces to the traditional $L_X \propto \sigma^4$ scaling law from simpler, dimensional arguments (e.g., Quintana & Melnick 1982). However, equation (9) tells us that c is a function of the velocity dispersion and the slope of the primordial spectrum. In fact, once $\tilde{\rho}(y)$ is specified, equations (7), (9), (15), and (16) may be used to eliminate V_{200} , c , $C(c)$, and $D(c)$, and one may thus obtain the dependence of L_X on σ_p and n . Specifically, suppose that within the range of interest for c , $C(c)$ and $D(c)$ exhibit a power law behavior of the form $C(c) \propto c^p$ and $D(c) \propto c^q$. Then

$$L_X \propto f^2 Z^{1/2+\xi/6} \sigma_p^{4-\xi}, \quad (23)$$

where

$$\xi = \frac{3(3+n)(3+3p-4q)}{3+14p-9q+n(6p-3-3q)} \quad (24)$$

2.3. Model Dependency

I consider dark matter density profiles of the form $\tilde{\rho}(y) = y^{-a}(1+y^b)^{-d}$. Thus each profile's shape may be specified by a set of three numbers, (a, b, d) . Some common profiles and their properties are listed in Table 1.

Once the set (a, b, d) is specified, $C(c)$ and $D(c)$ are readily computable. The relevant range of c for systems of galaxies comes from measurements of surface number density profiles and of r_{200} in clusters and groups of galaxies (Carlberg et al. 1997; Mahdavi et al. 1999). In these works, halos with masses in the range $10^{14} - 10^{16} M_\odot$ have concentrations $c \approx 2.5 - 9.5$, in good agreement with N-body simulations (e.g., NFW). For all sets (a, b, d) in Table 1, the power law approximations $C(c) \propto c^p$ and $D(c) \propto c^q$ are better than 8% accurate everywhere within $c = 2.5 - 9.5$.

Figure 1 shows $\xi(n)$ from equation (24) for various profiles. In all cases, $\xi(n)$ is positive, and approaches zero as $n \rightarrow -3$. As $n \rightarrow 0$, the models all predict a significant flattening of the $L_X - \sigma_p$ relation. This is understandable through equation (8): the characteristic density δ_c is highly anticorrelated with halo mass in the $n \approx 0$ universes, and hence the emission measure increases quite slowly or not at all with mass. As $n \rightarrow -3$, δ_c becomes nearly independent of the halo mass, and therefore L_X increases rapidly with the velocity dispersion.

3. APPLICATION

For low-redshift clusters of galaxies, $Z(z, \Omega) \approx 1$ with high accuracy. The dependence of the gas mass fraction f on $\langle T \rangle$ and hence σ_p is poorly determined. Some, using standard analysis, claim it should increase slightly with $\langle T \rangle$ (e.g., David, Jones, & Forman 1995); others say that accounting for cooling flows should cause it to decrease slightly with $\langle T \rangle$ (e.g. Allen & Fabian 1998). Mohr, Mathiesen, & Evrard (1999), in their detailed study of clusters with ROSAT pointings, find that within 1 Mpc f is nearly independent of $\langle T \rangle$. Because of the range of contrasting findings, I adopt $f \propto \langle T \rangle^{0.0 \pm 0.2} \propto \sigma_p^{0.0 \pm 0.4}$ at the 68% confidence level.

There is more agreement among observers regarding the empirical value of α_3 : Quintana & Melnick (1982) have $\alpha_3 = 4.0 \pm 0.7$ for data from the Einstein satellite; theirs are 2-10 keV luminosities, which scale similarly to bolometric luminosities for rich clusters. More recently, Mulchaey & Zabludoff (1998; MZ98) combined ROSAT observations of groups and clusters with deep optical spectroscopy to obtain $\alpha_3 = 4.29 \pm 0.37$ for bolometric luminosities. I adopt the average value, $\alpha_3 = 4.15 \pm 0.4$.

Assuming normally distributed errors, the adopted values of f and α_3 constrain ξ to be < 1.0 at the one-sided 90% confidence level, or < 1.93 at the one-sided 99% confidence level. Figure (1) shows the $\xi = 1.0, 1.93$ boundaries. To accommodate all the models I consider in this scenario, n must be < -2.0 at 90% confidence level, and < -1.1 at the 99% confidence level.

At least two systematic effects could bias these results. Preheating of the plasma in $k\langle T \rangle \lesssim 4$ keV clusters (Ponman, Cannon, & Navarro 1999) might suppress their luminosities, while leaving those of $k\langle T \rangle \gtrsim 4$ keV clusters unchanged. One can avoid this bias by ignoring all clusters with $k\langle T \rangle < 4$ keV. I find that removing these clusters, which make up $\approx 25\%$ of the MZ98 sample, does not sig-

nificantly affect the slope of the observed $L_X - \sigma$ relation.

Also, cooling flows might affect the luminosities. While Markevitch (1998) finds that the slope of the $L_X - \langle T \rangle$ relation does not change as a result of removing cooling flows, Allen & Fabian (1998) find that $\alpha_2 = 3.1 \pm 0.6$ changes to $\alpha_2 = 2.3 \pm 0.4$ after including cooling flows. The Markevitch (1998) method, which removes the cooling flow altogether, is more appropriate for probing the gravitational potential than the Allen & Fabian (1998) method, which includes the luminosity of the cooling component.

To constrain the effect of cooling flows on α_3 , I conduct the following test. Of the MZ98 clusters, 34% are also contained in Markevitch (1998). I fit α_3 for just these clusters, using the L_X which excludes the cooling component. I find that the slope changes to $\alpha_3 = 3.85 \pm 0.3$; with this, the upper limits on n become $n < -1.7$ and $n < -0.9$ at the one-sided 90% and 99% confidence levels. The chief result of this paper—that if n were much greater than -1 , α_3 should be ≈ 2 instead of the observed value, ≈ 4 —is therefore not affected by the 10% correction due to cooling flows.




FIG. 1.— The correction to the slope of the $L_X - \sigma_p$ relation, $\xi(n)$, for various density profiles. The dashed lines represent the one-sided 90% and 99% upper bounds on ξ from available observations. The arrows indicate the same confidence intervals on n .

4. CONCLUSION

If the characteristic densities of clusters of galaxies trace the background density of the universe at the time of each cluster's formation, there should be a relatively simple relationship between the x-ray luminosity L_X , the observed velocity dispersion σ_p , and the slope of the primordial power spectrum, n . This relationship, given by equations (23) and (24), depends slightly on the density profile of the clustered dark matter, but should not significantly depend on Ω or Λ when applied to low-redshift clusters of galaxies. For a wide range of assumed density profiles, the observations imply $n < -2.0$ and $n < -1.1$ at the one-sided 90% and 99% confidence levels, respectively. This is consistent with the bounds from the cluster temperature

TABLE 1
DENSITY PROFILES CONSIDERED

Model	a, b, d	p	q
Hernquist (1990)	1,1,3	0.35	0.16
NFW	1,1,2	0.74	0.58
Plummer Sphere	0,2,5/2	0.14	0.05
King (1962)	0,2,3/2	0.74	0.59
Jaffe (1983)	2,1,2	0.17	0.03

function, which give n between -2.3 and -1.15 . Improving this constraint depends largely on a better understanding of preheating, cooling flows, and the variation of the gas mass fraction with σ_p .

I thank the referee, Vincent Eke, for comments which improved the paper. I am grateful to Margaret Geller for our many discussions. Conversations with Saurabh Jha were also useful. This work was supported by the Smithsonian Institution and by the National Science Foundation.

REFERENCES

- Allen, S. W., & Fabian, A. C. 1998, MNRAS, 297, L57
 Binney, J., and Tremaine, S. 1987, Galactic Dynamics (Princeton: Princeton University Press)
 Carlberg, R. G., et al. 1997, ApJ, 495, L13
 David, L. P., Jones, C., & Forman, W. 1995, ApJ, 445, 578
 Eke, V., Navarro, J. F., & Frenk, C. S. 1998, ApJ, 503, 569
 Henry, J. P. & Arnaud, K. A. 1991, ApJ, 372, 410
 Henry, J. P. 1997, ApJ, 489, L1
 Hernquist, L. 1990, ApJ, 356, 359
 Jaffe, W. 1983, MNRAS, 202, 995
 King, I. R. 1962, AJ, 67, 471
 Mahdavi, A., Geller, M. J., Böhringer, H., Kurtz, M. J., & Ramella, M. 1997, ApJ, in press (astro-ph/9901095)
 Markevitch, M. 1998, ApJ, 504, 27
 Mohr, J. J., Mathiesen, N., & Evrard, A. E. 1999, ApJ, in press (astro-ph/9901281)
 Mulchaey, J. S., & Zabludoff, A. I. 1998, ApJ, 496, 73
 Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, ApJ, 490, 493
 Ponman, T. J., Cannon, D. B., & Navarro, J. F. 1999, Nature, 397, 135
 Quintana, H., and Melnick, J., 1982, AJ, 87, 972